

Student: _____
Date: _____

Instructor: Alfredo Alvarez
Course: 2413 Cal I

Assignment:
CAL2413ANSWERSI150FIESTA

1. The function $s(t)$ represents the position of an object at time t moving along a line. Suppose $s(1) = 120$ and $s(5) = 220$. Find the average velocity of the object over the interval of time $[1, 5]$.

The average velocity over the interval $[1, 5]$ is $v_{av} =$. (Simplify your answer.)

Answer: 25

ID: 2.1.3

2. The position of an object moving along a line is given by the function $s(t) = -8t^2 + 64t$. Find the average velocity of the object over the following intervals.

(a) $[1, 7]$

(b) $[1, 6]$

(c) $[1, 5]$

(d) $[1, 1 + h]$ where $h > 0$ is any real number.

(a) The average velocity of the object over the interval $[1, 7]$ is .

(b) The average velocity of the object over the interval $[1, 6]$ is .

(c) The average velocity of the object over the interval $[1, 5]$ is .

(d) The average velocity of the object over the interval $[1, 1 + h]$ is .

Answers 0

8

16

$-8h + 48$

ID: 2.1.13

3. For the position function $s(t) = -16t^2 + 106t$, complete the following table with the appropriate average velocities. Then make a conjecture about the value of the instantaneous velocity at $t = 1$.

Time Interval	[1, 2]	[1, 1.5]	[1, 1.1]	[1, 1.01]	[1, 1.001]
Average Velocity	_____	_____	_____	_____	_____

Complete the following table.

Time Interval	[1, 2]	[1, 1.5]	[1, 1.1]	[1, 1.01]	[1, 1.001]
Average Velocity	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>

(Type exact answers. Type integers or decimals.)

The value of the instantaneous velocity at $t = 1$ is .

(Round to the nearest integer as needed.)

Answers 58

66

72.4

73.84

73.984

74

ID: 2.1.17

4. For the function $f(x) = 12x^3 - x$, make a table of slopes of secant lines and make a conjecture about the slope of the tangent line at $x = 1$.

Complete the table.

(Round the final answer to three decimal places as needed. Round all intermediate values to four decimal places as needed.)

Interval	Slope of secant line
[1, 2]	<input type="text"/>
[1, 1.5]	<input type="text"/>
[1, 1.1]	<input type="text"/>
[1, 1.01]	<input type="text"/>
[1, 1.001]	<input type="text"/>

An accurate conjecture for the slope of the tangent line at $x = 1$ is .

(Round to the nearest integer as needed.)

Answers 83.000

56.000

38.720

35.360

35.000

35

ID: 2.1.28

5.

Let $f(x) = \frac{x^2 - 4}{x - 2}$. **(a)** Calculate $f(x)$ for each value of x in the following table. **(b)** Make a conjecture about the value of

$$\lim_{x \rightarrow 2} \frac{x^2 - 4}{x - 2}.$$

(a) Calculate $f(x)$ for each value of x in the following table.

x	1.9	1.99	1.999	1.9999
$f(x) = \frac{x^2 - 4}{x - 2}$				
x	2.1	2.01	2.001	2.0001
$f(x) = \frac{x^2 - 4}{x - 2}$				

(Type an integer or decimal rounded to four decimal places as needed.)

(b) Make a conjecture about the value of $\lim_{x \rightarrow 2} \frac{x^2 - 4}{x - 2}$.

$$\lim_{x \rightarrow 2} \frac{x^2 - 4}{x - 2} = \text{[]} \text{ (Type an integer or a decimal.)}$$

Answers 3.9

3.99

3.999

3.9999

4.1

4.01

4.001

4.0001

4

ID: 2.2.7

6. Let $g(t) = \frac{t - 64}{\sqrt{t} - 8}$.

a. Make two tables, one showing the values of g for $t = 63.9, 63.99, \text{ and } 63.999$ and one showing values of g for $t = 64.1, 64.01, \text{ and } 64.001$.

b. Make a conjecture about the value of $\lim_{t \rightarrow 64} \frac{t - 64}{\sqrt{t} - 8}$.

a. Make a table showing the values of g for $t = 63.9, 63.99, \text{ and } 63.999$.

t	63.9	63.99	63.999
g(t)			

(Round to four decimal places.)

Make a table showing the values of g for $t = 64.1, 64.01, \text{ and } 64.001$.

t	64.1	64.01	64.001
g(t)			

(Round to four decimal places.)

b. Make a conjecture about the value of $\lim_{t \rightarrow 64} \frac{t - 64}{\sqrt{t} - 8}$. Select the correct choice below and fill in any answer boxes in your choice.

☐ A. $\lim_{t \rightarrow 64} \frac{t - 64}{\sqrt{t} - 8} =$ _____ (Simplify your answer.)

☐ B. The limit does not exist.

Answers 15.9937

15.9994

15.9999

16.0062


16.0006

16.0001

A. $\lim_{t \rightarrow 64} \frac{t - 64}{\sqrt{t} - 8} =$ (Simplify your answer.)

ID: 2.2.9

7. Use the graph to find the following limits and function value.



- a. $\lim_{x \rightarrow 0^-} f(x)$
- b. $\lim_{x \rightarrow 0^+} f(x)$
- c. $\lim_{x \rightarrow 0} f(x)$
- d. $f(0)$

a. Find the limit. Select the correct choice below and fill in any answer boxes in your choice.

☐ A. $\lim_{x \rightarrow 0^-} f(x) =$ _____ (Type an integer.)

☐ B. The limit does not exist.

b. Find the limit. Select the correct choice below and fill in any answer boxes in your choice.

☐ A. $\lim_{x \rightarrow 0^+} f(x) =$ _____ (Type an integer.)

☐ B. The limit does not exist.

c. Find the limit. Select the correct choice below and fill in any answer boxes in your choice.

☐ A. $\lim_{x \rightarrow 0} f(x) =$ _____ (Type an integer.)

☐ B. The limit does not exist.

d. Find the function value. Select the correct choice below and fill in any answer boxes in your choice.

☐ A. $f(0) =$ _____ (Type an integer.)

☐ B. The answer is undefined.

Answers A. $\lim_{x \rightarrow 0^-} f(x) =$ (Type an integer.)

A. $\lim_{x \rightarrow 0^+} f(x) =$ (Type an integer.)

B. The limit does not exist.

A. $f(0) =$ (Type an integer.)

ID: 2.2.15

8. Explain why $\lim_{x \rightarrow -5} \frac{x^2 - 2x - 35}{x + 5} = \lim_{x \rightarrow -5} (x - 7)$, and then evaluate $\lim_{x \rightarrow -5} \frac{x^2 - 2x - 35}{x + 5}$.

Choose the correct answer below.

- ☐ A. The numerator of the expression $\frac{x^2 - 2x - 35}{x + 5}$ simplifies to $x - 7$ for all x , so the limits are equal.
- ☐ B. Since $\frac{x^2 - 2x - 35}{x + 5} = x - 7$ whenever $x \neq -5$, it follows that the two expressions evaluate to the same number as x approaches -5 .
- ☐ C. Since each limit approaches -5 , it follows that the limits are equal.
- ☐ D. The limits $\lim_{x \rightarrow -5} \frac{x^2 - 2x - 35}{x + 5}$ and $\lim_{x \rightarrow -5} (x - 7)$ equal the same number when evaluated using direct substitution.

Now evaluate the limit.

$$\lim_{x \rightarrow -5} \frac{x^2 - 2x - 35}{x + 5} = \boxed{} \text{ (Simplify your answer.)}$$

Answers B.

Since $\frac{x^2 - 2x - 35}{x + 5} = x - 7$ whenever $x \neq -5$, it follows that the two expressions evaluate to the same number as x approaches -5 .

-12

ID: 2.3.5

9. Assume $\lim_{x \rightarrow 9} f(x) = 9$ and $\lim_{x \rightarrow 9} h(x) = 3$. Compute the following limit and state the limit laws used to justify the computation.

$$\lim_{x \rightarrow 9} \frac{f(x)}{h(x)}$$

$$\lim_{x \rightarrow 9} \frac{f(x)}{h(x)} = \boxed{} \text{ (Simplify your answer.)}$$

Select each limit law used to justify the computation.

- ☐ A. Quotient
☐ B. Difference
☐ C. Root
☐ D. Sum
☐ E. Constant multiple
☐ F. Power
☐ G. Product

Answers 3

A. Quotient

ID: 2.3.8

10. Find the following limit or state that it does not exist.

$$\lim_{x \rightarrow 441} \frac{\sqrt{x} - 21}{x - 441}$$

Simplify the given limit.

$$\lim_{x \rightarrow 441} \frac{\sqrt{x} - 21}{x - 441} = \lim_{x \rightarrow 441} \left(\boxed{} \right) \text{ (Simplify your answer.)}$$

Evaluate the limit, if possible. Select the correct choice below and, if necessary, fill in the answer box to complete your choice.

- ☐ A. $\lim_{x \rightarrow 441} \frac{\sqrt{x} - 21}{x - 441} = \underline{\hspace{2cm}}$ (Type an exact answer.)
☐ B. The limit does not exist.

Answers $\frac{1}{\sqrt{x} + 21}$

A. $\lim_{x \rightarrow 441} \frac{\sqrt{x} - 21}{x - 441} = \boxed{\frac{1}{42}} \text{ (Type an exact answer.)}$

ID: 2.3.41-Setup & Solve

11. Determine the following limit at infinity.

$$\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} \text{ if } f(x) \rightarrow 900,000 \text{ and } g(x) \rightarrow \infty \text{ as } x \rightarrow \infty$$

Find the limit. Choose the correct answer below.

- ☐ A. $\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = 0$
- ☐ B. $\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = \infty$
- ☐ C. $\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = 900,000$
- ☐ D. $\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = \frac{1}{900,000}$

Answer: A. $\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = 0$

ID: 2.5.11

12. Determine the following limit at infinity.

$$\lim_{x \rightarrow \infty} \frac{8 + 9x + 7x^3}{x^3}$$

Select the correct choice below and, if necessary, fill in the answer box to complete your choice.

- ☐ A. $\lim_{x \rightarrow \infty} \frac{8 + 9x + 7x^3}{x^3} = \underline{\hspace{2cm}}$
- ☐ B. The limit does not exist and is neither $-\infty$ nor ∞ .

Answer: A. $\lim_{x \rightarrow \infty} \frac{8 + 9x + 7x^3}{x^3} =$

ID: 2.5.12

13. Determine the following limit.

$$\lim_{w \rightarrow \infty} \frac{10w^2 + 5w + 3}{\sqrt{4w^4 + w^3}}$$

Select the correct choice below, and, if necessary, fill in the answer box to complete your choice.

- ☐ A. $\lim_{w \rightarrow \infty} \frac{10w^2 + 5w + 3}{\sqrt{4w^4 + w^3}} = \underline{\hspace{2cm}}$ (Simplify your answer.)
- ☐ B. The limit does not exist and is neither ∞ nor $-\infty$.

Answer: A. $\lim_{w \rightarrow \infty} \frac{10w^2 + 5w + 3}{\sqrt{4w^4 + w^3}} = \boxed{5}$ (Simplify your answer.)

ID: 2.5.29

14. Determine the following limit.

$$\lim_{x \rightarrow -\infty} \frac{\sqrt{169x^2 + x}}{x}$$

Select the correct choice below, and, if necessary, fill in the answer box to complete your choice.

- ☐ A. $\lim_{x \rightarrow -\infty} \frac{\sqrt{169x^2 + x}}{x} = \underline{\hspace{2cm}}$ (Simplify your answer.)
- ☐ B. The limit does not exist and is neither ∞ nor $-\infty$.

Answer: A. $\lim_{x \rightarrow -\infty} \frac{\sqrt{169x^2 + x}}{x} = \boxed{-13}$ (Simplify your answer.)

ID: 2.5.31

15. Determine $\lim_{x \rightarrow \infty} f(x)$ and $\lim_{x \rightarrow -\infty} f(x)$ for the following function. Then give the horizontal asymptotes of f , if any.

$$f(x) = \frac{4x}{8x + 5}$$

Evaluate $\lim_{x \rightarrow \infty} f(x)$. Select the correct choice below and, if necessary, fill in the answer box to complete your choice.

☐ A. $\lim_{x \rightarrow \infty} \frac{4x}{8x + 5} =$ _____ (Simplify your answer.)

☐ B. The limit does not exist and is neither ∞ nor $-\infty$.

Evaluate $\lim_{x \rightarrow -\infty} f(x)$. Select the correct choice below and, if necessary, fill in the answer box to complete your choice.

☐ A. $\lim_{x \rightarrow -\infty} \frac{4x}{8x + 5} =$ _____ (Simplify your answer.)

☐ B. The limit does not exist and is neither ∞ nor $-\infty$.

Give the horizontal asymptotes of f , if any. Select the correct choice below and, if necessary, fill in the answer box(es) to complete your choice.

☐ A. The function has one horizontal asymptote, _____.
(Type an equation.)

☐ B. The function has two horizontal asymptotes. The top asymptote is _____ and the bottom asymptote is _____.
(Type equations.)

☐ C. The function has no horizontal asymptotes.

Answers A. $\lim_{x \rightarrow \infty} \frac{4x}{8x + 5} =$ (Simplify your answer.)

A. $\lim_{x \rightarrow -\infty} \frac{4x}{8x + 5} =$ (Simplify your answer.)

A. The function has one horizontal asymptote, . (Type an equation.)

ID: 2.5.37

16. Evaluate $\lim_{x \rightarrow \infty} f(x)$ and $\lim_{x \rightarrow -\infty} f(x)$ for the following rational function. Use ∞ or $-\infty$ where appropriate. Then give the horizontal asymptote of f (if any).

$$f(x) = \frac{4x^2 - 7x + 8}{2x^2 + 3}$$

Select the correct choice below and, if necessary, fill in the answer box to complete your choice.

- ☐ A. $\lim_{x \rightarrow \infty} f(x) =$ _____ (Simplify your answer.)
- ☐ B. The limit does not exist and is neither ∞ nor $-\infty$.

Select the correct choice below and, if necessary, fill in the answer box to complete your choice.

- ☐ A. $\lim_{x \rightarrow -\infty} f(x) =$ _____ (Simplify your answer.)
- ☐ B. The limit does not exist and is neither ∞ nor $-\infty$.

Identify the horizontal asymptote. Select the correct choice below and, if necessary, fill in the answer box to complete your choice.

- ☐ A. The horizontal asymptote is $y =$ _____.
- ☐ B. There are no horizontal asymptotes.

Answers A. $\lim_{x \rightarrow \infty} f(x) =$ (Simplify your answer.)

A. $\lim_{x \rightarrow -\infty} f(x) =$ (Simplify your answer.)

A. The horizontal asymptote is $y =$.

ID: 2.5.39

17. Evaluate $\lim_{x \rightarrow \infty} f(x)$ and $\lim_{x \rightarrow -\infty} f(x)$ for the rational function. Then give the horizontal asymptote of f (if any).

$$f(x) = \frac{20x^6 - 8}{4x^6 - 9x^5}$$

Select the correct choice below, and, if necessary, fill in the answer box to complete your choice.

☐ A. $\lim_{x \rightarrow -\infty} \frac{20x^6 - 8}{4x^6 - 9x^5} = \underline{\hspace{2cm}}$ (Simplify your answer.)

- ☐ B. The limit does not exist and is neither ∞ or $-\infty$.

Select the correct choice below, and, if necessary, fill in the answer box to complete your choice.

☐ A. $\lim_{x \rightarrow \infty} \frac{20x^6 - 8}{4x^6 - 9x^5} = \underline{\hspace{2cm}}$ (Simplify your answer.)

- ☐ B. The limit does not exist and is neither ∞ or $-\infty$.

Select the correct choice below, and, if necessary, fill in the answer box to complete your choice.

- ☐ A. The function has a horizontal asymptote at $y = \underline{\hspace{2cm}}$.
(Simplify your answer.)

- ☐ B. The function has no horizontal asymptote.

Answers A. $\lim_{x \rightarrow -\infty} \frac{20x^6 - 8}{4x^6 - 9x^5} = \boxed{5}$ (Simplify your answer.)

A. $\lim_{x \rightarrow \infty} \frac{20x^6 - 8}{4x^6 - 9x^5} = \boxed{5}$ (Simplify your answer.)

A. The function has a horizontal asymptote at $y = \boxed{5}$. (Simplify your answer.)

ID: 2.5.40

18. Determine $\lim_{x \rightarrow \infty} f(x)$ and $\lim_{x \rightarrow -\infty} f(x)$ for the following function. Then give the horizontal asymptotes of f , if any.

$$f(x) = \frac{3x^3 - 7}{x^4 + 5x^2}$$

Evaluate $\lim_{x \rightarrow \infty} f(x)$. Select the correct choice below and, if necessary, fill in the answer box to complete your choice.

☐ A. $\lim_{x \rightarrow \infty} \frac{3x^3 - 7}{x^4 + 5x^2} = \underline{\hspace{2cm}}$ (Simplify your answer.)

☐ B. The limit does not exist and is neither ∞ nor $-\infty$.

Evaluate $\lim_{x \rightarrow -\infty} f(x)$. Select the correct choice below and, if necessary, fill in the answer box to complete your choice.

☐ A. $\lim_{x \rightarrow -\infty} \frac{3x^3 - 7}{x^4 + 5x^2} = \underline{\hspace{2cm}}$ (Simplify your answer.)

☐ B. The limit does not exist and is neither ∞ nor $-\infty$.

Identify the horizontal asymptotes. Select the correct choice below and, if necessary, fill in the answer box(es) to complete your choice.

☐ A. The function has one horizontal asymptote, $\underline{\hspace{2cm}}$.
(Type an equation.)

☐ B. The function has two horizontal asymptotes. The top asymptote is $\underline{\hspace{2cm}}$ and the bottom asymptote is $\underline{\hspace{2cm}}$.
(Type equations.)

☐ C. The function has no horizontal asymptotes.

Answers A. $\lim_{x \rightarrow \infty} \frac{3x^3 - 7}{x^4 + 5x^2} = \boxed{0}$ (Simplify your answer.)

A. $\lim_{x \rightarrow -\infty} \frac{3x^3 - 7}{x^4 + 5x^2} = \boxed{0}$ (Simplify your answer.)

A. The function has one horizontal asymptote, $\boxed{y = 0}$. (Type an equation.)

ID: 2.5.41

19. Find all the asymptotes of the function.

$$f(x) = \frac{8x^2 + 24}{2x^2 + 3x - 2}$$

Find the horizontal asymptotes. Select the correct choice below and, if necessary, fill in the answer box(es) to complete your choice.

- ☐ A. The function has one horizontal asymptote, _____.
(Type an equation using y as the variable.)
- ☐ B. The function has two horizontal asymptotes. The top asymptote is _____ and the bottom asymptote is _____.
(Type equations using y as the variable.)
- ☐ C. The function has no horizontal asymptotes.

Find the vertical asymptotes. Select the correct choice below and, if necessary, fill in the answer box(es) to complete your choice.

- ☐ A. The function has one vertical asymptote, _____.
(Type an equation using x as the variable.)
- ☐ B. The function has two vertical asymptotes. The leftmost asymptote is _____ and the rightmost asymptote is _____.
(Type equations using x as the variable.)
- ☐ C. The function has no vertical asymptotes.

Find the slant asymptote(s). Select the correct choice below and, if necessary, fill in the answer box(es) to complete your choice.

- ☐ A. The function has one slant asymptote, _____.
(Type an equation using x and y as the variables.)
- ☐ B. The function has two slant asymptotes. The asymptote with the larger slope is _____ and the asymptote with the smaller slope is _____.
(Type equations using x and y as the variables.)
- ☐ C. The function has no slant asymptotes.

Answers A. The function has one horizontal asymptote, . (Type an equation using y as the variable.)

B.

The function has two vertical asymptotes. The leftmost asymptote is and the rightmost

asymptote is .

(Type equations using x as the variable.)

C. The function has no slant asymptotes.

ID: EXTRA 2.66

20. Determine whether the following function is continuous at a . Use the continuity checklist to justify your answer.

$$f(x) = \frac{4x^2 + 25x + 25}{x^2 - 8x}, a = 8$$

Select all that apply.

- ☐ A. The function is continuous at $a = 8$.
- ☐ B. The function is not continuous at $a = 8$ because $f(8)$ is undefined.
- ☐ C. The function is not continuous at $a = 8$ because $\lim_{x \rightarrow 8} f(x)$ does not exist.
- ☐ D. The function is not continuous at $a = 8$ because $\lim_{x \rightarrow 8} f(x) \neq f(8)$.

Answer: B. The function is not continuous at $a = 8$ because $f(8)$ is undefined. , C.

The function is not continuous at $a = 8$ because $\lim_{x \rightarrow 8} f(x)$ does not exist. , D.

The function is not continuous at $a = 8$ because $\lim_{x \rightarrow 8} f(x) \neq f(8)$.

ID: 2.6.17

21. Determine whether the following function is continuous at a . Use the continuity checklist to justify your answer.

$$f(x) = \begin{cases} \frac{x^2 - 144}{x - 12} & \text{if } x \neq 12 \\ 3 & \text{if } x = 12 \end{cases}; a = 12$$

Select all that apply.

- ☐ A. The function is continuous at $a = 12$.
- ☐ B. The function is not continuous at $a = 12$ because $f(12)$ is undefined.
- ☐ C. The function is not continuous at $a = 12$ because $\lim_{x \rightarrow 12} f(x)$ does not exist.
- ☐ D. The function is not continuous at $a = 12$ because $\lim_{x \rightarrow 12} f(x) \neq f(12)$.

Answer: D. The function is not continuous at $a = 12$ because $\lim_{x \rightarrow 12} f(x) \neq f(12)$.

ID: 2.6.21

22. Determine the intervals on which the following function is continuous.

$$f(x) = \frac{x^2 - 5x + 6}{x^2 - 4}$$

On what interval(s) is f continuous?

(Simplify your answer. Type your answer in interval notation. Use a comma to separate answers as needed.)

Answer: $(-\infty, -2), (-2, 2), (2, \infty)$

ID: 2.6.28

23. Evaluate the following limit.

$$\lim_{x \rightarrow 5} \sqrt{x^2 + 24}$$

Select the correct choice below and, if necessary, fill in the answer box to complete your choice.

- ☒ A. $\lim_{x \rightarrow 5} \sqrt{x^2 + 24} =$, because $x^2 + 24$ is continuous for all x and the square root function is continuous for all $x \geq 0$.
(Type an integer or a fraction.)
- ☐ B. The limit does not exist and is neither ∞ nor $-\infty$.

Answer: A.

$\lim_{x \rightarrow 5} \sqrt{x^2 + 24} =$, because $x^2 + 24$ is continuous for all x and the square root function is continuous for all $x \geq 0$.
(Type an integer or a fraction.)

ID: 2.6.53

24. Suppose x lies in the interval $(1, 3)$ with $x \neq 2$. Find the smallest positive value of δ such that the inequality $0 < |x - 2| < \delta$ is true for all possible values of x .

The smallest positive value of δ is . (Type an integer or a fraction.)

Answer: 1

ID: 2.7.1

25. Suppose $|f(x) - 7| < 0.3$ whenever $0 < x < 7$. Find all values of $\delta > 0$ such that $|f(x) - 7| < 0.3$ whenever $0 < |x - 2| < \delta$.

The values of δ are $0 < \delta \leq$. (Type an integer or a fraction.)

Answer: 2

ID: 2.7.7


26. The function f in the figure satisfies $\lim_{x \rightarrow 2} f(x) = 4$. Determine the largest value of $\delta > 0$ satisfying each statement.

a. If $0 < |x - 2| < \delta$, then $|f(x) - 4| < 2$.

b. If $0 < |x - 2| < \delta$, then $|f(x) - 4| < 1$.

a. $\delta =$ (Simplify your answer.)

b. $\delta =$ (Simplify your answer.)



Answers 1

$\frac{1}{2}$

ID: 2.7.9

27. Use the precise definition of a limit to prove the following limit. Specify a relationship between ε and δ that guarantees the limit exists.

$$\lim_{x \rightarrow 0} (6x + 9) = 9$$

Let $\varepsilon > 0$ be given. Choose the correct proof below.

- ☐ A. Choose $\delta = \varepsilon$. Then, $|(6x + 9) - 9| < \varepsilon$ whenever $0 < |x - 0| < \delta$.
- ☐ B. Choose $\delta = 6\varepsilon$. Then, $|(6x + 9) - 9| < \varepsilon$ whenever $0 < |x - 0| < \delta$.
- ☐ C. Choose $\delta = \frac{\varepsilon}{9}$. Then, $|(6x + 9) - 9| < \varepsilon$ whenever $0 < |x - 0| < \delta$.
- ☐ D. Choose $\delta = \frac{\varepsilon}{6}$. Then, $|(6x + 9) - 9| < \varepsilon$ whenever $0 < |x - 0| < \delta$.
- ☐ E. None of the above proofs is correct.

Answer:

D. Choose $\delta = \frac{\varepsilon}{6}$. Then, $|(6x + 9) - 9| < \varepsilon$ whenever $0 < |x - 0| < \delta$.

ID: 2.7.19

28. Find the average velocity of the function over the given interval.

$$y = \frac{3}{x-2}, [4, 7]$$

- ☐ A. 2
- ☐ B. 7
- ☐ C. $\frac{1}{3}$
- ☐ D. $-\frac{3}{10}$

Answer:

D. $-\frac{3}{10}$

ID: 2.1-3

29. Find all vertical asymptotes of the given function.

$$g(x) = \frac{x + 11}{x^2 + 16x}$$

- ☐ A. $x = 0, x = -16$
☐ B. $x = -16, x = -11$
☐ C. $x = 0, x = -4, x = 4$
☐ D. $x = -4, x = 4$

Answer: A. $x = 0, x = -16$

ID: 2.4-19

30. Divide numerator and denominator by the highest power of x in the denominator to find the limit at infinity.

$$\lim_{x \rightarrow \infty} \frac{\sqrt[3]{x} - 3x - 5}{-6x + x^{2/3} - 7}$$

- ☐ A. $\frac{1}{2}$
☐ B. $-\infty$
☐ C. 2
☐ D. 0

Answer: A. $\frac{1}{2}$


ID: 2.5-12

31. Use the graph to find a $\delta > 0$ such that for all x , $0 < |x - x_0| < \delta$ and $|f(x) - L| < \varepsilon$. Use the following information:
 $f(x) = 2x + 3$, $\varepsilon = 0.2$, $x_0 = 1$, $L = 5$.

¹ Click the icon to view the graph.

- ☐ A. 4
☐ B. 0.2
☐ C. 0.4
☐ D. 0.1

1: Graph



Answer: D. 0.1

ID: 2.7-1

32. Find the value of the derivative of the function at the given point.

$$f(x) = 4x^2 - 5x; (-1, 9)$$

$f'(-1) =$ (Type an integer or a simplified fraction.)

Answer: -13

ID: 3.1.1

33. a. Use the definition $m_{\tan} = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$ to find the slope of the line tangent to the graph of f at P .
 b. Determine an equation of the tangent line at P .

$$f(x) = x^2 - 3, P(-5, 22)$$

a. $m_{\tan} =$


b. $y =$

Answers - 10

- 10x - 28


ID: 3.1.25

34. Match the graph of the function on the right with the graph of its derivative.




Choose the correct graph of the function (in blue) and its derivative (in red) below.


☐ A.




☐ B.




☐ C.



☐ D.



Answer:



D.

ID: 3.2.49

35. A line perpendicular to another line or to a tangent line is called a normal line. Find an equation of the line perpendicular to the line that is tangent to the following curve at the given point P.

$$y = 7x - 15; P(2, -1)$$

The equation of the normal line at P(2, -1) is .

Answer: $y = -\frac{1}{7}x + \left(-\frac{5}{7}\right)$

ID: 3.2.63

36. Evaluate the derivative of the function given below using a limit definition of the derivative.

$$f(x) = x^2 + 6x - 8$$

$f'(x) =$

Answer: $2x + 6$

ID: EXTRA 3.2

37. Use the Quotient Rule to evaluate and simplify $\frac{d}{dx} \left(\frac{x-4}{2x-5} \right)$.

$\frac{d}{dx} \left(\frac{x-4}{2x-5} \right) =$

Answer: $\frac{3}{(2x-5)^2}$

ID: 3.4.5

38. Use the Quotient Rule to find $g'(1)$ given that $g(x) = \frac{2x^2}{6x+1}$.

$g'(1) =$
(Simplify your answer.)

Answer: $\frac{16}{49}$

ID: 3.4.6

39. **a.** Use the Product Rule to find the derivative of the given function.
b. Find the derivative by expanding the product first.

$$f(x) = (x - 3)(4x + 2)$$

a. Use the product rule to find the derivative of the function. Select the correct answer below and fill in the answer box(es) to complete your choice.

- ☐ **A.** The derivative is $(x - 3)(\quad) + (4x + 2)(\quad)$.
☐ **B.** The derivative is $(x - 3)(4x + 2)(\quad)$.
☐ **C.** The derivative is $(\quad)(x - 3)$.
☐ **D.** The derivative is $(\quad)x(4x + 2)$.
☐ **E.** The derivative is $(x - 3)(4x + 2) + (\quad)$.

b. Expand the product.

$$(x - 3)(4x + 2) = \boxed{} \text{ (Simplify your answer.)}$$

$$\text{Using either approach, } \frac{d}{dx}(x - 3)(4x + 2) = \boxed{}.$$

$$\text{Answers A. The derivative is } (x - 3)(\boxed{4}) + (4x + 2)(\boxed{1}).$$

$$4x^2 - 10x - 6$$

$$8x - 10$$

ID: 3.4.9

40. Use the quotient rule to find the derivative of the given function. Then find the derivative by first simplifying the function. Are the results the same?

$$h(w) = \frac{5w^4 - w}{w}$$

What is the immediate result of applying the quotient rule? Select the correct answer below.

- ☐ A. $(20w^3 - 1)(w) + (5w^4 - w)(1)$
- ☐ B. $15w^2$
- ☐ C. $5w^3 - 1$
- ☐ D. $\frac{w(20w^3 - 1) - (5w^4 - w)(1)}{w^2}$

What is the fully simplified result of applying the quotient rule?

What is the result of first simplifying the function, then taking the derivative? Select the correct answer below.

- ☐ A. $15w^2$
- ☐ B. $5w^3 - 1$
- ☐ C. $(20w^3 - 1)(w) + (5w^4 - w)(1)$
- ☐ D. $\frac{w(20w^3 - 1) - (5w^4 - w)(1)}{w^2}$

Are the two results the same?

- ☐ Yes
- ☐ No

Answers

D. $\frac{w(20w^3 - 1) - (5w^4 - w)(1)}{w^2}$

$$15w^2$$

A. $15w^2$

Yes

ID: 3.4.11

41. If $f(x) = -\cos x$, then what is the value of $f'(\pi)$?

$f'(\pi) =$ (Simplify your answer.)

Answer: 0

ID: 3.5.5

42. Evaluate the limit.

$$\lim_{x \rightarrow 0} \frac{\sin 3x}{\sin 2x}$$

Select the correct choice below and, if necessary, fill in the answer box to complete your choice.

- ☐ A. $\lim_{x \rightarrow 0} \frac{\sin 3x}{\sin 2x} = \underline{\hspace{2cm}}$
- ☐ B. The limit is undefined.

Answer: A. $\lim_{x \rightarrow 0} \frac{\sin 3x}{\sin 2x} = \boxed{\frac{3}{2}}$

ID: 3.5.13

43. Find $\frac{dy}{dx}$ for the following function.

$$y = 7 \sin x + 9 \cos x$$

$$\frac{dy}{dx} = \boxed{\hspace{2cm}}$$

Answer: $7 \cos x - 9 \sin x$

ID: 3.5.23

44. Find the derivative of the following function.

$$y = e^{-x} \sin x$$

$$\frac{dy}{dx} = \boxed{\hspace{2cm}}$$

Answer: $e^{-x}(\cos x - \sin x)$

ID: 3.5.25

45. Find an equation of the line tangent to the following curve at the given point.

$$y = 2x^2 + 3 \sin x; P(0,0)$$

The equation for the tangent line is $\boxed{\hspace{2cm}}$.

Answer: $y = 3x$

ID: EXTRA 3.73

46.

Let $h(x) = f(g(x))$ and $p(x) = g(f(x))$. Use the table below to compute the following derivatives.

a. $h'(1)$ b. $p'(4)$

x	1	2	3	4
$f(x)$	3	2	4	1
$f'(x)$	-2	-5	-4	-6
$g(x)$	4	1	2	3
$g'(x)$	$\frac{1}{5}$	$\frac{2}{5}$	$\frac{3}{5}$	$\frac{4}{5}$

Answers $-\frac{6}{5}$

$-\frac{6}{5}$

ID: 3.7.25

47. Calculate the derivative of the following function.

$$y = (5x + 1)^7$$

$$\frac{dy}{dx} = \boxed{}$$

Answer: $35(5x + 1)^6$

ID: 3.7.27

48. Calculate the derivative of the following function.

$$y = 3(6x^5 + 7)^{-4}$$

$$\frac{dy}{dx} = \boxed{}$$

Answer: $\frac{-360x^4}{(6x^5 + 7)^5}$

ID: 3.7.31

$h'(1) = \boxed{}$ (Simplify your answer.)

$p'(4) = \boxed{}$ (Simplify your answer.)

49. Calculate the derivative of the following function.

$$y = \cos(3t + 20)$$

$$\frac{dy}{dt} = \boxed{}$$

Answer: $-3 \sin(3t + 20)$

ID: 3.7.32

50. Calculate the derivative of the following function.

$$y = \tan(e^x)$$

$$\frac{dy}{dx} = \boxed{}$$

Answer: $e^x \sec^2 e^x$

ID: 3.7.35

51. Calculate the derivative of the following function.

$$y = \sin(6 \cos x)$$

$$\frac{dy}{dx} = \boxed{}$$

Answer: $-6 \cos(6 \cos x) \cdot \sin x$

ID: 3.7.44

52. Calculate $\frac{dy}{dx}$ using implicit differentiation.

$$16x = y^4$$

$$\frac{dy}{dx} = \boxed{}$$

Answer: $\frac{4}{y^3}$

ID: 3.8.5

53. Calculate $\frac{dy}{dx}$ using implicit differentiation.

$$\cos(y) + 8 = x$$

$$\frac{dy}{dx} = \boxed{}$$

Answer: - **csc** y

ID: 3.8.7

54. Consider the curve $x = y^3$. Use implicit differentiation to verify that $\frac{dy}{dx} = \frac{1}{3y^2}$ and then find $\frac{d^2y}{dx^2}$.

Use implicit differentiation to find the derivative of each side of the equation.

$$\frac{d}{dx}x = \boxed{} \text{ and } \frac{d}{dx}y^3 = \boxed{} \frac{dy}{dx}$$

Solve for $\frac{dy}{dx}$.

$$\frac{dy}{dx} = \boxed{}$$

Find $\frac{d^2y}{dx^2}$.

$$\frac{d^2y}{dx^2} = \boxed{}$$

Answers 1

$$3y^2$$

$$\frac{1}{3y^2}$$

$$-\frac{2}{9y^5}$$

ID: 3.8.11

55. Carry out the following steps for the given curve.

a. Use implicit differentiation to find $\frac{dy}{dx}$.

b. Find the slope of the curve at the given point.

$$x^3 + y^3 = -63; (-4, 1)$$

a. Use implicit differentiation to find $\frac{dy}{dx}$.

$$\frac{dy}{dx} = \boxed{}$$

b. Find the slope of the curve at the given point.

The slope of $x^3 + y^3 = -63$ at $(-4, 1)$ is $\boxed{}$.
(Simplify your answer.)

Answers $-\frac{x^2}{y^2}$
- 16

ID: 3.8.13

56. Use implicit differentiation to find $\frac{dy}{dx}$.

$$\cos(y) + \sin(x) = 2y$$

$$\frac{dy}{dx} = \boxed{}$$

Answer: $\frac{\cos x}{2 + \sin y}$

ID: 3.8.27

57. Use implicit differentiation to find $\frac{dy}{dx}$.

$$3 \cos(xy) = 4x + 7y$$

$$\frac{dy}{dx} = \boxed{}$$

Answer: $\frac{4 + 3y \sin(xy)}{-3x \sin(xy) - 7}$

ID: 3.8.31

58. Use implicit differentiation to find $\frac{dy}{dx}$.

$$e^{3xy} = 8y$$

$$\frac{dy}{dx} = \boxed{}$$

Answer: $\frac{3y e^{3xy}}{8 - 3x e^{3xy}}$

ID: 3.8.32

59. Use implicit differentiation to find $\frac{dy}{dx}$ for the following equation.

$$5x^5 + 7y^5 = 12xy$$

$$\frac{dy}{dx} = \boxed{}$$

Answer: $\frac{25x^4 - 12y}{12x - 35y^4}$

ID: 3.8.37

60. Find $\frac{d}{dx} (\ln \sqrt{x^2 + 9})$.

$$\frac{d}{dx} (\ln \sqrt{x^2 + 9}) = \boxed{}$$

Answer: $\frac{x}{x^2 + 9}$

ID: 3.9.9

61. Express the function $f(x) = g(x)^{h(x)}$ in terms of the natural logarithmic and natural exponential functions (base e).

$$f(x) = \boxed{}$$

Answer: $e^{h(x) \ln g(x)}$

ID: 3.9.11

62. Find the derivative.

$$\frac{d}{dx} (\ln (2x^2 + 3))$$

$$\frac{d}{dx} (\ln (2x^2 + 3)) = \boxed{}$$

Answer: $\frac{4x}{2x^2 + 3}$

ID: 3.9.17

63. Evaluate the derivative.

$$y = 2x^{2\pi}$$

$$y' = \boxed{} \text{ (Type an exact answer.)}$$

Answer: $4\pi x^{(2\pi - 1)}$

ID: 3.9.33

64. Find $\frac{dy}{dx}$ for the function $y = 17^x$.

$$\frac{dy}{dx} = \boxed{}$$

Answer: $17^x \ln 17$

ID: 3.9.37

65. Calculate the derivative of the following function.

$$y = 8 \log_4 (x^3 - 3)$$

$$\frac{d}{dx} 8 \log_4 (x^3 - 3) = \boxed{}$$

Answer: $\frac{24x^2}{(x^3 - 3) \ln 4}$

ID: 3.9.63

66. Use logarithmic differentiation to evaluate $f'(x)$.

$$f(x) = \frac{(x+4)^8}{(2x-4)^{12}}$$

$f'(x) =$

Answer: $\frac{(x+4)^8}{(2x-4)^{12}} \left[\frac{8}{x+4} - \frac{12}{x-2} \right]$

ID: 3.9.77

67. State the derivative formulas for $\sin^{-1}x$, $\tan^{-1}x$, and $\sec^{-1}x$.

What is the derivative of $\sin^{-1}x$?

- ☐ A. $\frac{1}{|x|\sqrt{x^2-1}}$ for $|x| > 1$
- ☐ B. $-\frac{1}{|x|\sqrt{x^2-1}}$ for $|x| > 1$
- ☐ C. $\frac{1}{\sqrt{1-x^2}}$ for $-1 < x < 1$
- ☐ D. $-\frac{1}{\sqrt{1-x^2}}$ for $-1 < x < 1$

What is the derivative of $\tan^{-1}x$?

- ☐ A. $-\frac{1}{|x|\sqrt{x^2-1}}$ for $|x| > 1$
- ☐ B. $-\frac{1}{\sqrt{1-x^2}}$ for $-1 < x < 1$
- ☐ C. $-\frac{1}{1+x^2}$ for $-\infty < x < \infty$
- ☐ D. $\frac{1}{1+x^2}$ for $-\infty < x < \infty$

What is the derivative of $\sec^{-1}x$?

- ☐ A. $-\frac{1}{|x|\sqrt{x^2-1}}$ for $|x| > 1$
- ☐ B. $-\frac{1}{\sqrt{1-x^2}}$ for $-1 < x < 1$
- ☐ C. $\frac{1}{|x|\sqrt{x^2-1}}$ for $|x| > 1$
- ☐ D. $\frac{1}{\sqrt{1-x^2}}$ for $-1 < x < 1$

Answers C. $\frac{1}{\sqrt{1-x^2}}$ for $-1 < x < 1$

D. $\frac{1}{1+x^2}$ for $-\infty < x < \infty$

C. $\frac{1}{|x|\sqrt{x^2-1}}$ for $|x| > 1$

ID: 3.10.1

68. Evaluate the derivative of the function.

$$f(x) = \sin^{-1}(9x^4)$$

$$f'(x) = \boxed{}$$

Answer: $\frac{36x^3}{\sqrt{1-81x^8}}$

ID: 3.10.13

69. Find the derivative of the function $y = 2 \tan^{-1}(2x)$.

$$\frac{dy}{dx} = \boxed{}$$

Answer: $\frac{4}{1+(2x)^2}$

ID: 3.10.19

70. Evaluate the derivative of the following function.

$$f(s) = \cot^{-1}(e^s)$$

$$\frac{d}{ds} \cot^{-1}(e^s) = \boxed{}$$

Answer: $-\frac{e^s}{1+e^{2s}}$

ID: 3.10.39

71. The sides of a square increase in length at a rate of 3 m/sec.

- a. At what rate is the area of the square changing when the sides are 15 m long?
 b. At what rate is the area of the square changing when the sides are 28 m long?

a. Write an equation relating the area of a square, A , and the side length of the square, s .

Differentiate both sides of the equation with respect to t .

$$\frac{dA}{dt} = \left(\frac{\quad}{\quad} \right) \frac{ds}{dt}$$

The area of the square is changing at a rate of (1) when the sides are 15 m long.

b. The area of the square is changing at a rate of (2) when the sides are 28 m long.

- | | |
|-----------------------------------|-------------------------------|
| (1) <input type="radio"/> m^3/s | (2) <input type="radio"/> m |
| <input type="radio"/> m/s | <input type="radio"/> m/s |
| <input type="radio"/> m^2/s | <input type="radio"/> m^2/s |
| <input type="radio"/> m | <input type="radio"/> m^3/s |

Answers $A = s^2$

$2s$

90

(1) m^2/s

168

(2) m^2/s

ID: 3.11.11-Setup & Solve

72. The area of a circle increases at a rate of $5 \text{ cm}^2 / \text{s}$.

- a. How fast is the radius changing when the radius is 4 cm?
 b. How fast is the radius changing when the circumference is 4 cm?

a. Write an equation relating the area of a circle, A , and the radius of the circle, r .

(Type an exact answer, using π as needed.)

Differentiate both sides of the equation with respect to t .

$$\frac{dA}{dt} = \left(\frac{\quad}{\quad} \right) \frac{dr}{dt}$$

(Type an exact answer, using π as needed.)

When the radius is 4 cm, the radius is changing at a rate of (1)
 (Type an exact answer, using π as needed.)

b. When the circumference is 4 cm, the radius is changing at a rate of (2)
 (Type an exact answer, using π as needed.)

- (1) ☐ cm^3 / s . (2) ☐ cm .
☐ cm . ☐ cm^2 / s .
☐ cm / s . ☐ cm / s .
☐ cm^2 / s . ☐ cm^3 / s .

Answers $A = \pi r^2$

$$2\pi r$$

$$\frac{5}{8\pi}$$

(1) cm / s .

$$\frac{5}{4}$$

(2) cm / s .

ID: 3.11.15-Setup & Solve

73. The edges of a cube increase at a rate of 2 cm / s. How fast is the volume changing when the length of each edge is 30 cm?

Write an equation relating the volume of a cube, V , and an edge of the cube, a .

Differentiate both sides of the equation with respect to t .

$$\frac{dV}{dt} = \left(\text{ } \right) \frac{da}{dt} \text{ (Type an expression using } a \text{ as the variable.)}$$

The rate of change of the volume is (1)
(Simplify your answer.)

- (1) ☐ cm / sec. ☐ cm^3 .
☐ cm^2 / sec. ☐ cm.
☐ cm^3 / sec.
☐ cm^2 .

Answers $V = a^3$

$$3a^2$$

$$5400$$

$$(1) \text{ cm}^3 / \text{sec.}$$

ID: 3.11.16-Setup & Solve

74. Find an equation for the tangent to the curve at the given point.

$$y = x^2 - 1, (-2, 3)$$

- ☐ A. $y = -2x - 5$
☐ B. $y = -4x - 9$
☐ C. $y = -4x - 10$
☐ D. $y = -4x - 5$

Answer: D. $y = -4x - 5$

ID: 3.1-2

75. At time t , the position of a body moving along the s -axis is $s = t^3 - 12t^2 + 36t$ m. Find the displacement of the body from $t = 0$ to $t = 3$.

- ☐ A. 27 m
☐ B. 63 m
☐ C. 37 m
☐ D. 32 m

Answer: A. 27 m

ID: 3.6-3

76. Use implicit differentiation to find dy/dx .

$$xy + x = 2$$

- ☐ A. $-\frac{1+y}{x}$
☐ B. $\frac{1+x}{y}$
☐ C. $-\frac{1+x}{y}$
☐ D. $\frac{1+y}{x}$

Answer: A. $-\frac{1+y}{x}$

ID: 3.8-4

77. Find the derivative of the function.

$$y = \log_2 \sqrt{9x + 4}$$

- ☐ A. $\frac{9}{\ln 2}$
- ☐ B. $\frac{9}{\ln 2 (9x + 4)}$
- ☐ C. $\frac{9}{2(\ln 2)(9x + 4)}$
- ☐ D. $\frac{9 \ln 2}{9x + 4}$

Answer: C. $\frac{9}{2(\ln 2)(9x + 4)}$

ID: 3.9-11


78. Boyle's law states that if the temperature of a gas remains constant, then $PV = c$, where P = pressure, V = volume, and c is a constant. Given a quantity of gas at constant temperature, if V is decreasing at a rate of $10 \text{ in}^3/\text{sec}$, at what rate is P increasing when $P = 50 \text{ lb/in}^2$ and $V = 90 \text{ in}^3$? (Do not round your answer.)

- ☐ A. $\frac{50}{9} \text{ lb / in}^2 \text{ per sec}$
- ☐ B. $450 \text{ lb / in}^2 \text{ per sec}$
- ☐ C. $\frac{25}{81} \text{ lb / in}^2 \text{ per sec}$
- ☐ D. $18 \text{ lb / in}^2 \text{ per sec}$

Answer: A. $\frac{50}{9} \text{ lb / in}^2 \text{ per sec}$

ID: 3.11-7

79. Determine from the graph whether the function has any absolute extreme values on $[a, b]$.



Where do the absolute extreme values of the function occur on $[a, b]$?

- ☐ A. There is no absolute maximum and the absolute minimum occurs at $x = a$ on $[a, b]$.
☐ B. There is no absolute maximum and there is no absolute minimum on $[a, b]$.
☐ C. The absolute maximum occurs at $x = d$ and the absolute minimum occurs at $x = a$ on $[a, b]$.
☐ D. The absolute maximum occurs at $x = d$ and there is no absolute minimum on $[a, b]$.

Answer: C. The absolute maximum occurs at $x = d$ and the absolute minimum occurs at $x = a$ on $[a, b]$.

ID: 4.1.13

80. Find the critical points of the following function.

$$f(x) = 2x^2 - 3x + 1$$

What is the derivative of $f(x) = 2x^2 - 3x + 1$?

$$f'(x) = \boxed{}$$

Find the critical points, if any, of f on the domain. Select the correct choice below and, if necessary, fill in the answer box to complete your choice.

- ☐ A. The critical point(s) occur(s) at $x = $.
 (Use a comma to separate answers as needed.)
☐ B. There are no critical points for $f(x) = 2x^2 - 3x + 1$ on the domain.

Answers $4x - 3$

A. The critical point(s) occur(s) at $x = \boxed{\frac{3}{4}}$. (Use a comma to separate answers as needed.)

ID: 4.1.23-Setup & Solve

81. Find the critical points of the following function.

$$f(x) = -\frac{x^3}{3} + 9x$$

Select the correct choice below and, if necessary, fill in the answer box to complete your choice.

- ☐ A. The critical point(s) occur(s) at $x =$ _____.
(Use a comma to separate answers as needed.)
- ☐ B. There are no critical points.

Answer: A. The critical point(s) occur(s) at $x =$. (Use a comma to separate answers as needed.)

ID: 4.1.25

82. Determine the location and value of the absolute extreme values of f on the given interval, if they exist.

$$f(x) = -x^2 + 8 \text{ on } [-3, 4]$$

What is/are the absolute maximum/maxima of f on the given interval? Select the correct choice below and, if necessary, fill in the answer boxes to complete your choice.

- ☐ A. The absolute maximum/maxima is/are _____ at $x =$ _____.
(Use a comma to separate answers as needed.)
- ☐ B. There is no absolute maximum of f on the given interval.

What is/are the absolute minimum/minima of f on the given interval? Select the correct choice below and, if necessary, fill in the answer boxes to complete your choice.

- ☐ A. The absolute minimum/minima is/are _____ at $x =$ _____.
(Use a comma to separate answers as needed.)
- ☐ B. There is no absolute minimum of f on the given interval.

Answers A. The absolute maximum/maxima is/are at $x =$.

(Use a comma to separate answers as needed.)

A. The absolute minimum/minima is/are at $x =$.

(Use a comma to separate answers as needed.)

ID: 4.1.43

83. A stone is launched vertically upward from a cliff 336 ft above the ground at a speed of 64 ft/s. Its height above the ground t seconds after the launch is given by $s = -16t^2 + 64t + 336$ for $0 \leq t \leq 7$. When does the stone reach its maximum height?

Find the derivative of s .

$$s' = \boxed{}$$

The stone reaches its maximum height at $\boxed{}$ s.
(Simplify your answer.)

Answers $-32t + 64$
2

ID: 4.1.73

84. Suppose a tour guide has a bus that holds a maximum of 84 people. Assume his profit (in dollars) for taking n people on a city tour is $P(n) = n(42 - 0.5n) - 84$. (Although P is defined only for positive integers, treat it as a continuous function.)

- a. How many people should the guide take on a tour to maximize the profit?
b. Suppose the bus holds a maximum of 35 people. How many people should be taken on a tour to maximize the profit?

a. Find the derivative of the given function $P(n)$.

$$P'(n) = \boxed{}$$

If the bus holds a maximum of 84 people, the guide should take $\boxed{}$ people on a tour to maximize the profit.

b. If the bus holds a maximum of 35 people, the guide should take $\boxed{}$ people on a tour to maximize the profit.

Answers $-n + 42$
42
35

ID: 4.1.75

85. At what points c does the conclusion of the Mean Value Theorem hold for $f(x) = x^3$ on the interval $[-17, 17]$?

The conclusion of the Mean Value Theorem holds for $c = \boxed{}$.

(Use a comma to separate answers as needed. Type an exact answer, using radicals as needed.)

$$\text{Answer: } \frac{17\sqrt{3}}{3}, -\frac{17\sqrt{3}}{3}$$

ID: 4.2.8

86. **a.** Determine whether the Mean Value Theorem applies to the function $f(x) = -1 + x^2$ on the interval $[-2, 1]$.
b. If so, find the point(s) that are guaranteed to exist by the Mean Value Theorem.
-

a. Choose the correct answer below.

- ☐ **A.** Yes, because the function is continuous on the interval $[-2, 1]$ and differentiable on the interval $(-2, 1)$.
☐ **B.** No, because the function is differentiable on the interval $(-2, 1)$, but is not continuous on the interval $[-2, 1]$.
☐ **C.** No, because the function is not continuous on the interval $[-2, 1]$, and is not differentiable on the interval $(-2, 1)$.
☐ **D.** No, because the function is continuous on the interval $[-2, 1]$, but is not differentiable on the interval $(-2, 1)$.

b. Select the correct choice below and, if necessary, fill in the answer box to complete your choice.

- ☐ **A.** The point(s) is/are $x =$ _____.
(Simplify your answer. Use a comma to separate answers as needed.)
☐ **B.** The Mean Value Theorem does not apply in this case.

Answers **A.** Yes, because the function is continuous on the interval $[-2, 1]$ and differentiable on the interval $(-2, 1)$.

A. The point(s) is/are $x =$.


(Simplify your answer. Use a comma to separate answers as needed.)

ID: 4.2.21




87. Sketch a function that is continuous on $(-\infty, \infty)$ and has the following properties. Use a number line to summarize information about the function.

$$f'(x) < 0 \text{ on } (-\infty, 3); f'(x) > 0 \text{ on } (3, 7); f'(x) < 0 \text{ on } (7, \infty).$$

Which number line summarizes the information about the function?



Which of the following graphs matches the description of the given properties?



ID: 4.3.9

88. Find the intervals on which f is increasing and the intervals on which it is decreasing.

$$f(x) = -7 + x^2$$

Select the correct choice below and, if necessary, fill in the answer box(es) to complete your choice.

- ☐ A. The function is increasing on _____ and decreasing on _____.
(Simplify your answers. Type your answers in interval notation. Use a comma to separate answers as needed.)
- ☐ B. The function is increasing on _____. The function is never decreasing.
(Simplify your answer. Type your answer in interval notation. Use a comma to separate answers as needed.)
- ☐ C. The function is decreasing on _____. The function is never increasing.
(Simplify your answer. Type your answer in interval notation. Use a comma to separate answers as needed.)
- ☐ D. The function is never increasing nor decreasing.

Answer: A. The function is increasing on and decreasing on .

(Simplify your answers. Type your answers in interval notation. Use a comma to separate answers as needed.)

ID: 4.3.19

89. Find the intervals on which f is increasing and the intervals on which it is decreasing.

$$f(x) = 4 - 3x + 2x^2$$

Select the correct choice below and, if necessary, fill in the answer box(es) to complete your choice.

- ☐ A. The function is increasing on _____ and decreasing on _____.
(Simplify your answers. Type your answers in interval notation. Use a comma to separate answers as needed.)
- ☐ B. The function is increasing on _____. The function is never decreasing.
(Simplify your answer. Type your answer in interval notation. Use a comma to separate answers as needed.)
- ☐ C. The function is decreasing on _____. The function is never increasing.
(Simplify your answer. Type your answer in interval notation. Use a comma to separate answers as needed.)
- ☐ D. The function is never increasing nor decreasing.

Answer: A. The function is increasing on and decreasing on .

(Simplify your answers. Type your answers in interval notation. Use a comma to separate answers as needed.)

ID: 4.3.25

90. Locate the critical points of the following function. Then use the Second Derivative Test to determine whether they correspond to local maxima, local minima, or neither.

$$f(x) = x^3 - 15x^2$$

What is(are) the critical point(s) of f ? Select the correct choice below and, if necessary, fill in the answer box to complete your choice.

- ☐ A. The critical point(s) is(are) $x =$ _____.
(Use a comma to separate answers as needed. Type an integer or a simplified fraction.)
- ☐ B. There are no critical points for f .

Find $f''(x)$.

$$f''(x) = \boxed{}$$

What is/are the local maximum/maxima of f ? Select the correct choice below and, if necessary, fill in the answer box to complete your choice.

- ☐ A. The local maximum/maxima of f is/are at $x =$ _____.
(Use a comma to separate answers as needed. Type an integer or a simplified fraction.)
- ☐ B. There is no local maximum of f .

What is/are the local minimum/minima of f ? Select the correct choice below and, if necessary, fill in the answer box to complete your choice.

- ☐ A. The local minimum/minima of f is/are at $x =$ _____.
(Use a comma to separate answers as needed. Type an integer or a simplified fraction.)
- ☐ B. There is no local minimum of f .

Answers A. The critical point(s) is(are) $x = \boxed{0,10}$.
(Use a comma to separate answers as needed. Type an integer or a simplified fraction.)

$$6x - 30$$

A. The local maximum/maxima of f is/are at $x = \boxed{0}$.
(Use a comma to separate answers as needed. Type an integer or a simplified fraction.)

A. The local minimum/minima of f is/are at $x = \boxed{10}$.
(Use a comma to separate answers as needed. Type an integer or a simplified fraction.)

ID: 4.3.77-Setup & Solve

91. Locate the critical points of the following function. Then use the Second Derivative Test to determine whether they correspond to local maxima, local minima, or neither.

$$f(x) = 2 - 2x^2$$

What is(are) the critical point(s) of f ? Select the correct choice below and, if necessary, fill in the answer box to complete your choice.

- ☐ A. The critical point(s) is(are) $x =$ _____.
(Use a comma to separate answers as needed. Type an integer or a simplified fraction.)
- ☐ B. There are no critical points for f .

What is/are the local maximum/maxima of f ? Select the correct choice below and, if necessary, fill in the answer box to complete your choice.

- ☐ A. The local maximum/maxima of f is/are at $x =$ _____.
(Use a comma to separate answers as needed. Type an integer or a simplified fraction.)
- ☐ B. There is no local maximum of f .

What is/are the local minimum/minima of f ? Select the correct choice below and, if necessary, fill in the answer box to complete your choice.

- ☐ A. The local minimum/minima of f is/are at $x =$ _____.
(Use a comma to separate answers as needed. Type an integer or a simplified fraction.)
- ☐ B. There is no local minimum of f .

Answers A. The critical point(s) is(are) $x =$.
(Use a comma to separate answers as needed. Type an integer or a simplified fraction.)

A. The local maximum/maxima of f is/are at $x =$.
(Use a comma to separate answers as needed. Type an integer or a simplified fraction.)

B. There is no local minimum of f .

ID: 4.3.79

92. Locate the critical points of the following function. Then use the Second Derivative Test to determine whether they correspond to local maxima, local minima, or neither.

$$f(x) = -3x^3 + 9x^2 + 4$$

What is(are) the critical point(s) of f ? Select the correct choice below and, if necessary, fill in the answer box to complete your choice.

- ☐ A. The critical point(s) is(are) $x =$ _____.
(Use a comma to separate answers as needed. Type an integer or a simplified fraction.)
- ☐ B. There are no critical points for f .

What is/are the local minimum/minima of f ? Select the correct choice below and, if necessary, fill in the answer box to complete your choice.

- ☐ A. The local minimum/minima of f is/are at $x =$ _____.
(Use a comma to separate answers as needed. Type an integer or a simplified fraction.)
- ☐ B. There is no local minimum of f .

What is/are the local maximum/maxima of f ? Select the correct choice below and, if necessary, fill in the answer box to complete your choice.

- ☐ A. The local maximum/maxima of f is/are at $x =$ _____.
(Use a comma to separate answers as needed. Type an integer or a simplified fraction.)
- ☐ B. There is no local maximum of f .

Answers A. The critical point(s) is(are) $x =$.
(Use a comma to separate answers as needed. Type an integer or a simplified fraction.)

A. The local minimum/minima of f is/are at $x =$.
(Use a comma to separate answers as needed. Type an integer or a simplified fraction.)

A. The local maximum/maxima of f is/are at $x =$.
(Use a comma to separate answers as needed. Type an integer or a simplified fraction.)

ID: 4.3.83

93. **a.** Squares with sides of length x are cut out of each corner of a rectangular piece of cardboard measuring 5 ft by 4 ft. The resulting piece of cardboard is then folded into a box without a lid. Find the volume of the largest box that can be formed in this way.
- b.** Suppose that in part (a) the original piece of cardboard is a square with sides of length s . Find the volume of the largest box that can be formed in this way.

a. To find the objective function, express the volume V of the box in terms of x .

$$V = \boxed{}$$

(Type an expression.)

The interval of interest of the objective function is $\boxed{}$.

(Simplify your answer. Type your answer in interval notation. Use integers or decimals for any numbers in the expression.)

The maximum volume of the box is approximately $\boxed{}$ ft^3 .

(Round to the nearest hundredth as needed.)

b. To find the objective function, express the volume V of the box in terms of s and x .

$$V = \boxed{}$$

(Type an expression.)

The maximum volume of the box is $\boxed{}$.

(Type an expression using s as the variable.)

Answers $x(5 - 2x)(4 - 2x)$

$[0, 2]$

6.56

$x(s - 2x)(s - 2x)$

$\frac{2s^3}{27}$

ID: 4.5.40

94. Use a linear approximation to estimate the following quantity. Choose a value of a to produce a small error.

$\ln(0.95)$

What is the value found using the linear approximation?

$\ln(0.95) \approx \boxed{}$ (Round to two decimal places as needed.)

Answer: -0.05

ID: 4.6.41

95. Consider the following function and express the relationship between a small change in x and the corresponding change in y in the form $dy = f'(x)dx$.

$$f(x) = 2x^3 - 5x$$

$$dy = (\text{ }) dx$$

$$\text{Answer: } 6x^2 - 5$$

ID: 4.6.67

96. Consider the following function and express the relationship between a small change in x and the corresponding change in y in the form $dy = f'(x) dx$.

$$f(x) = \cot 7x$$

$$dy = (\text{ }) dx$$

$$\text{Answer: } -7 \csc^2(7x)$$

ID: 4.6.69

97. Evaluate the following limit. Use l'Hôpital's Rule when it is convenient and applicable.

$$\lim_{x \rightarrow 0} \frac{5 \sin 2x}{3x}$$

Use l'Hôpital's Rule to rewrite the given limit so that it is not an indeterminate form.

$$\lim_{x \rightarrow 0} \frac{5 \sin 2x}{3x} = \lim_{x \rightarrow 0} (\text{ })$$

Evaluate the limit.

$$\lim_{x \rightarrow 0} \frac{5 \sin 2x}{3x} = (\text{ }) \text{ (Type an exact answer.)}$$

$$\text{Answers } \frac{10 \cos(2x)}{3}$$

$$\frac{10}{3}$$

ID: 4.7.29-Setup & Solve

98. Use a calculator or program to compute the first 10 iterations of Newton's method when they are applied to the following function with the given initial approximation.

$$f(x) = x^2 - 12; x_0 = 4$$

k	x_k	k	x_k
0	<input type="text"/>	6	<input type="text"/>
1	<input type="text"/>	7	<input type="text"/>
2	<input type="text"/>	8	<input type="text"/>
3	<input type="text"/>	9	<input type="text"/>
4	<input type="text"/>	10	<input type="text"/>
5	<input type="text"/>		

(Round to six decimal places as needed.)

Answers 4.000000

3.464102

3.500000

3.464102

3.464286

3.464102

3.464102

3.464102

3.464102

3.464102

3.464102

ID: 4.8.13-T

99. Use a calculator or program to compute the first 10 iterations of Newton's method for the given function and initial approximation.

$$f(x) = 4 \sin x + x + 1, x_0 = 1.2$$

Complete the table.

(Do not round until the final answer. Then round to six decimal places as needed.)

k	x_k	k	x_k
1	<input type="text"/>	6	<input type="text"/>
2	<input type="text"/>	7	<input type="text"/>
3	<input type="text"/>	8	<input type="text"/>
4	<input type="text"/>	9	<input type="text"/>
5	<input type="text"/>	10	<input type="text"/>

Answers – 1.220218

– 0.201082

0.455144

– 0.201082

– 0.244542

– 0.201082

– 0.200905

– 0.201082

– 0.201082

– 0.201082

ID: 4.8.15-T

100. Determine the following indefinite integral. Check your work by differentiation.

$$\int (9x^{17} - 5x^9) \, dx$$

$$\int (9x^{17} - 5x^9) \, dx = \boxed{} \text{ (Use } C \text{ as the arbitrary constant.)}$$

Answer: $\frac{x^{18}}{2} - \frac{x^{10}}{2} + C$

ID: 4.9.23

101. Evaluate the following indefinite integral.

$$\int \left(\frac{3}{\sqrt{x}} + 3\sqrt{x} \right) dx$$

$$\int \left(\frac{3}{\sqrt{x}} + 3\sqrt{x} \right) dx = \boxed{}$$

(Use C as the arbitrary constant.)

Answer:

$$6\sqrt{x} + 2x^{\frac{3}{2}} + C$$

ID: 4.9.25

102. Find $\int (6x + 5)^2 dx$.

$$\int (6x + 5)^2 dx = \boxed{}$$

(Use C as the arbitrary constant.)

Answer: $12x^3 + 30x^2 + 25x + C$

ID: 4.9.27

103. Determine the following indefinite integral. Check your work by differentiation.

$$\int 4m(12m^2 - 7m) dm$$

$$\int 4m(12m^2 - 7m) dm = \boxed{} \text{ (Use C as the arbitrary constant.)}$$

Answer:

$$12m^4 - \frac{28m^3}{3} + C$$

ID: 4.9.28

104. Determine the following indefinite integral. Check your work by differentiation.

$$\int \left(3x^{\frac{1}{3}} + 4x^{-\frac{2}{3}} + 9 \right) dx$$

$$\int \left(3x^{\frac{1}{3}} + 4x^{-\frac{2}{3}} + 9 \right) dx = \boxed{} \text{ (Use C as the arbitrary constant.)}$$

Answer: $\frac{9}{4}x^{\frac{4}{3}} + 12x^{\frac{1}{3}} + 9x + C$

ID: 4.9.29

105. Determine the following indefinite integral. Check your work by differentiation.

$$\int 3\sqrt[4]{x} \, dx$$

$$\int 3\sqrt[4]{x} \, dx = \boxed{} \text{ (Use C as the arbitrary constant.)}$$

Answer: $\frac{12}{5}x^{\frac{5}{4}} + C$

ID: 4.9.30

106. Determine the following indefinite integral. Check your work by differentiation.

$$\int (5x + 9)(1 - x) \, dx$$

$$\int (5x + 9)(1 - x) \, dx = \boxed{} \\ \text{(Use C as the arbitrary constant.)}$$

Answer: $-\frac{5}{3}x^3 - 2x^2 + 9x + C$

ID: 4.9.31

107. Determine the following indefinite integral.

$$\int \frac{3x^5 - 6x^4}{x^3} dx$$

$$\int \frac{3x^5 - 6x^4}{x^3} dx = \boxed{}$$

(Use C as the arbitrary constant.)

Answer: $x^3 - 3x^2 + C$

ID: 4.9.35

108. For the following function f, find the antiderivative F that satisfies the given condition.

$$f(x) = 2x^3 + 3 \sin x, F(0) = 3$$

The antiderivative that satisfies the given condition is $F(x) = \boxed{}$.

Answer: $\frac{1}{2}x^4 - 3 \cos x + 6$

ID: 4.9.71

109. For the following function f, find the antiderivative F that satisfies the given condition.

$$f(u) = 2e^u + 5; F(0) = 8$$

The antiderivative that satisfies the given condition is $F(u) = \boxed{}$.

Answer: $2e^u + 5u + 6$

ID: 4.9.74

110. Given the following velocity function of an object moving along a line, find the position function with the given initial position.

$$v(t) = 6t^2 + 4t - 7; s(0) = 0$$

The position function is $s(t) = \boxed{}$.

Answer: $2t^3 + 2t^2 - 7t$

ID: 4.9.95

111. Find the absolute extreme values of the function on the interval.

$$F(x) = \sqrt[3]{x}, \quad -8 \leq x \leq 27$$

- ☐ A. absolute maximum is 3 at $x = -27$; absolute minimum is -2 at $x = 27$
- ☐ B. absolute maximum is 3 at $x = 27$; absolute minimum is -2 at $x = -8$
- ☐ C. absolute maximum is 0 at $x = 0$; absolute minimum is -2 at $x = -8$
- ☐ D. absolute maximum is 3 at $x = 27$; absolute minimum is 0 at $x = 0$

Answer: B. absolute maximum is 3 at $x = 27$; absolute minimum is -2 at $x = -8$

ID: 4.1-13

112. Find the value or values of c that satisfy the equation $\frac{f(b) - f(a)}{b - a} = f'(c)$ in the conclusion of the Mean Value Theorem for the function and interval.

$$f(x) = x^2 + 3x + 3, \quad [-2, 3]$$


- ☐ A. $-\frac{1}{2}, \frac{1}{2}$
- ☐ B. $0, \frac{1}{2}$
- ☐ C. $\frac{1}{2}$
- ☐ D. $-2, 3$

Answer: C. $\frac{1}{2}$

ID: 4.2-1

113.

Use the graph of the function $f(x)$ to locate the local extrema and identify the intervals where the function is concave up and concave down.



- ☐ A. Local minimum at $x = 1$; local maximum at $x = -1$; concave down on $(-\infty, \infty)$
- ☐ B. Local minimum at $x = 1$; local maximum at $x = -1$; concave down on $(0, \infty)$; concave up on $(-\infty, 0)$
- ☐ C. Local minimum at $x = 1$; local maximum at $x = -1$; concave up on $(0, \infty)$; concave down on $(-\infty, 0)$
- ☐ D. Local minimum at $x = 1$; local maximum at $x = -1$; concave up on $(-\infty, \infty)$

Answer: C. Local minimum at $x = 1$; local maximum at $x = -1$; concave up on $(0, \infty)$; concave down on $(-\infty, 0)$

ID: 4.3-9

114. From a thin piece of cardboard 20 in by 20 in, square corners are cut out so that the sides can be folded up to make a box. What dimensions will yield a box of maximum volume? What is the maximum volume? Round to the nearest tenth, if necessary.

- ☐ A. 13.3 in \times 13.3 in \times 6.7 in; 1,185.2 in³
- ☐ B. 13.3 in \times 13.3 in \times 3.3 in; 592.6 in³
- ☐ C. 6.7 in \times 6.7 in \times 6.7 in; 296.3 in³
- ☐ D. 10.0 in \times 10.0 in \times 5.0 in; 500.0 in³

Answer: B. 13.3 in \times 13.3 in \times 3.3 in; 592.6 in³

ID: 4.5-1

115. Solve the initial value problem.

$$\frac{ds}{dt} = \cos t - \sin t, \quad s\left(\frac{\pi}{2}\right) = 7$$

- ☐ A. $s = \sin t + \cos t + 8$
- ☐ B. $s = 2 \sin t + 5$
- ☐ C. $s = \sin t - \cos t + 6$
- ☐ D. $s = \sin t + \cos t + 6$


Answer: D. $s = \sin t + \cos t + 6$

ID: 4.9-16


116. Suppose an object moves along a line at 18 m/s for $0 \leq t \leq 3$ s and at 25 m/s for $3 < t \leq 5$ s. Sketch the graph of the velocity function and find the displacement of the object for $0 \leq t \leq 5$.

Sketch the graph of the velocity function. Choose the correct graph below.


☐ A.




☐ B.



☐ C.




☐ D.



The displacement of the object for $0 \leq t \leq 5$ is m. (Simplify your answer.)

Answers




B.

104

ID: 5.1.1


117.

Approximate the area of the region bounded by the graph of $f(x)$ (shown below) and the x -axis by dividing the interval $[5, 9]$ into $n = 4$ subintervals. Use a left and right Riemann sum to obtain two different approximations. Draw the approximating rectangles.




In which graph below are the selected points the left endpoints of the 4 approximating rectangles?


☐ A.




☐ B.



☐ C.




☐ D.




Using the specified rectangles, approximate the area.

In which graph below are the selected points the right endpoints of the 4 approximating rectangles?


☐ A.




☐ B.



☐ C.




☐ D.




Using the specified rectangles, approximate the area.

Answers



D.

16



D.

13

ID: 5.1.9

118. Evaluate the following expressions.

a. $\sum_{k=1}^{16} k$

b. $\sum_{k=1}^5 (3k+2)$

c. $\sum_{k=1}^8 k^2$

d. $\sum_{n=1}^5 (3+n^2)$

e. $\sum_{m=1}^4 \frac{4m+4}{9}$

f. $\sum_{j=1}^3 (5j-6)$

g. $\sum_{k=1}^9 k(6k+5)$

h. $\sum_{n=0}^7 \sin \frac{n\pi}{2}$

a. $\sum_{k=1}^{16} k =$ (Type an integer or a simplified fraction.)

b. $\sum_{k=1}^5 (3k+2) =$ (Type an integer or a simplified fraction.)

c. $\sum_{k=1}^8 k^2 =$ (Type an integer or a simplified fraction.)

d. $\sum_{n=1}^5 (3+n^2) =$ (Type an integer or a simplified fraction.)

e. $\sum_{m=1}^4 \frac{4m+4}{9} =$ (Type an integer or a simplified fraction.)

f. $\sum_{j=1}^3 (5j-6) =$ (Type an integer or a simplified fraction.)

g. $\sum_{k=1}^9 k(6k+5) =$ (Type an integer or a simplified fraction.)

h. $\sum_{n=0}^7 \sin \frac{n\pi}{2} =$ (Type an integer or a simplified fraction.)

Answers 136

55

204

70

$\frac{56}{9}$

12

1935

0

ID: 5.1.49

119.

The functions f and g are integrable and $\int_3^7 f(x)dx = 6$, $\int_3^7 g(x)dx = 3$, and $\int_4^7 f(x)dx = 4$. Evaluate the integral below or state that there is not enough information.

$$-\int_7^3 3f(x)dx$$

Select the correct choice below and, if necessary, fill in the answer box to complete your choice.

☐ A. $-\int_7^3 3f(x)dx =$ _____ (Simplify your answer.)

☐ B. There is not enough information to evaluate $-\int_7^3 3f(x)dx$.

Answer: A. $-\int_7^3 3f(x)dx =$ (Simplify your answer.)

ID: EXTRA 5.18

120.

Evaluate $\frac{d}{dx} \int_a^x f(t) dt$ and $\frac{d}{dx} \int_a^b f(t) dt$, where a and b are constants.

$$\frac{d}{dx} \int_a^x f(t) dt =$$
 (Simplify your answer.)

$$\frac{d}{dx} \int_a^b f(t) dt =$$
 (Simplify your answer.)


Answers $f(x)$

0

ID: 5.3.9

121. Evaluate the following integral using the Fundamental Theorem of Calculus. Discuss whether your result is consistent with the figure.

$$\int_0^1 (x^2 - 3x + 4) dx$$



$$\int_0^1 (x^2 - 3x + 4) dx = \boxed{}$$

Is your result consistent with the figure?

- ☐ A. No, because the definite integral is positive and the graph of f lies below the x -axis.
- ☐ B. Yes, because the definite integral is positive and the graph of f lies above the x -axis.
- ☐ C. Yes, because the definite integral is negative and the graph of f lies below the x -axis.
- ☐ D. No, because the definite integral is negative and the graph of f lies above the x -axis.


Answers $\frac{17}{6}$

B. Yes, because the definite integral is positive and the graph of f lies above the x -axis.

ID: 5.3.23

122.

Evaluate the integral $\int_{-\pi/4}^{7\pi/4} (\sin x + \cos x) dx$ using the fundamental theorem of calculus. Discuss whether your result is consistent with the figure shown to the right.



$$\int_{-\pi/4}^{7\pi/4} (\sin x + \cos x) dx = \boxed{}$$

Is this value consistent with the given figure?

- ☐ A. The value is consistent with the figure because the total area can be approximated using a rectangle of base π and height $\sin(\pi/4) + \cos(\pi/4) = \sqrt{2}$.
- ☐ B. The value is not consistent with the figure because the total area could be approximated using a rectangle of base π and height $\sin(\pi/4) + \cos(\pi/4) = \sqrt{2}$.
- ☐ C. The value is not consistent with the figure because the figure is a graph of the base function, $f(x)$, instead of a graph of the area function, $A(x)$.
- ☐ D. The value is consistent with the figure because the area below the x-axis appears to be equal to the area above the x-axis.

Answers 0

D.

The value is consistent with the figure because the area below the x-axis appears to be equal to the area above the x-axis.

ID: 5.3.24


123. Evaluate the following integral using the fundamental theorem of calculus. Sketch the graph of the integrand and shade the region whose net area you have found.

$$\int_{-3}^1 (x^2 + 2x - 3) \, dx$$


$$\int_{-3}^1 (x^2 + 2x - 3) \, dx = \boxed{}$$

Choose the correct sketch below.


☐ A.




☐ B.




☐ C.



☐ D.



Answers $-\frac{32}{3}$



A.

ID: 5.3.25

124. Use the fundamental theorem of calculus to evaluate the following definite integral.

$$\int_0^3 (5x^2 + 4) \, dx$$

$$\int_0^3 (5x^2 + 4) \, dx = \boxed{}$$

(Type an exact answer.)

Answer: 57

ID: 5.3.29

125. Evaluate the following integral using the Fundamental Theorem of Calculus.

$$\int_4^9 (4-x)(x-9) \, dx$$

$$\int_4^9 (4-x)(x-9) \, dx = \boxed{}$$

(Type an exact answer.)

Answer: $\frac{125}{6}$

ID: 5.3.41

126. Find the area of the region bounded by the graph of f and the x -axis on the given interval.

$$f(x) = x^2 - 45; [3, 4]$$

The area is $\boxed{}$. (Type an integer or a simplified fraction.)

Answer: $\frac{98}{3}$

ID: 5.3.67

127. Simplify the following expression.

$$\frac{d}{dx} \int_x^3 \sqrt{t^2 + 1} \, dt$$

$$\frac{d}{dx} \int_x^3 \sqrt{t^2 + 1} \, dt = \boxed{}$$

Answer: $-\sqrt{x^2 + 1}$

ID: 5.3.75

128. Simplify the following expression.

$$\frac{d}{dx} \int_{10}^{x^3} \frac{dp}{p^2}$$

$$\frac{d}{dx} \int_{10}^{x^3} \frac{dp}{p^2} = \boxed{}$$

Answer: $\frac{3}{x^4}$

ID: 5.3.77

129. Evaluate the following integral.

$$\int_0^4 (x-5)^2 dx$$

$$\int_0^4 (x-5)^2 dx = \boxed{}$$

(Type an exact answer. Use C as the arbitrary constant as needed.)

Answer: $\frac{124}{3}$

ID: EXTRA 5.57


130. Find the average value of the following function over the given interval. Draw a graph of the function and indicate the average value.

$$f(x) = x(x - 1); [5, 7]$$


The average value of the function is $\bar{f} =$.

Choose the correct graph of $f(x)$ and \bar{f} below.


☐ A.




☐ B.




☐ C.



☐ D.



Answers $\frac{91}{3}$




ID: 5.4.30


131. The elevation of a path is given by $f(x) = x^3 - 3x^2 + 2$, where x measures horizontal distances. Draw a graph of the elevation function and find its average value for $0 \leq x \leq 4$.

Choose the correct graph below.


☐ A.




☐ B.



☐ C.




☐ D.



The average value is . (Type an integer or a simplified fraction.)

Answers



B.

2

ID: 5.4.34

132. Find the point(s) at which the function $f(x) = 5 - 6x$ equals its average value on the interval $[0, 4]$.

The function equals its average value at $x =$.

(Use a comma to separate answers as needed.)

Answer: 2

ID: 5.4.39

133. Use the substitution $u = x^2 + 15$ to find the following indefinite integral. Check your answer by differentiation.

$$\int 2x(x^2 + 15)^7 dx$$

$$\int 2x(x^2 + 15)^7 dx = \text{}$$

(Use C as the arbitrary constant.)

Answer: $\frac{1}{8}(x^2 + 15)^8 + C$

ID: 5.5.7

134. Use the substitution $u = 7x^2 - 1$ to find the following indefinite integral. Check your answer by differentiation.

$$\int 14x \cos(7x^2 - 1) \, dx$$

$$\int 14x \cos(7x^2 - 1) \, dx = \boxed{}$$

(Use C as the arbitrary constant.)

Answer: $\sin(7x^2 - 1) + C$

ID: 5.5.8

135. Use the substitution $u = 7x^2 + 4x$ to evaluate the indefinite integral below.

$$\int (14x + 4)\sqrt{7x^2 + 4x} \, dx$$

Write the integrand in terms of u .

$$\int (14x + 4)\sqrt{7x^2 + 4x} \, dx = \int (\boxed{}) \, du$$

Evaluate the integral.

$$\int (14x + 4)\sqrt{7x^2 + 4x} \, dx = \boxed{}$$

(Use C as the arbitrary constant.)

Answers \sqrt{u}

$$\frac{2}{3}(7x^2 + 4x)^{\frac{3}{2}} + C$$

ID: 5.5.10-Setup & Solve

136. Use a change of variables or the accompanying table to evaluate the following indefinite integral.

$$\int \frac{e^{2x}}{e^{2x} + 7} dx$$

² Click the icon to view the table of general integration formulas.

Determine a change of variables from x to u . Choose the correct answer below.

- ☐ A. $u = 2x$
- ☐ B. $u = e^{2x}$
- ☐ C. $u = e^{2x} + 7$
- ☐ D. $u = \frac{1}{e^{2x} + 7}$

Write the integral in terms of u .

$$\int \frac{e^{2x}}{e^{2x} + 7} dx = \int (\text{ }) du$$

Evaluate the integral.

$$\int \frac{e^{2x}}{e^{2x} + 7} dx = \text{ }$$

(Use C as the arbitrary constant.)

2: General Integration Formulas

$$\int \cos ax \, dx = \frac{1}{a} \sin ax + C$$

$$\int \sin ax \, dx = -\frac{1}{a} \cos ax + C$$

$$\int \sec^2 ax \, dx = \frac{1}{a} \tan ax + C$$

$$\int \csc^2 ax \, dx = -\frac{1}{a} \cot ax + C$$

$$\int \sec ax \tan ax \, dx = \frac{1}{a} \sec ax + C$$

$$\int \csc ax \cot ax \, dx = -\frac{1}{a} \csc ax + C$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax} + C$$

$$\int b^x dx = \frac{1}{\ln b} b^x + C, b > 0, b \neq 1$$

$$\int \frac{dx}{a^2 + x^2} = \frac{1}{a} \tan^{-1} \frac{x}{a} + C$$

$$\int \frac{dx}{\sqrt{a^2 - x^2}} = \sin^{-1} \frac{x}{a} + C, a > 0$$

$$\int \frac{dx}{x\sqrt{x^2 - a^2}} = \frac{1}{a} \sec^{-1} \left| \frac{x}{a} \right| + C, a > 0$$

Answers C. $u = e^{2x} + 7$

$$\frac{1}{2u}$$

$$\frac{1}{2} \ln |e^{2x} + 7| + C$$

ID: 5.5.44-Setup & Solve

137. Use a change of variables or the table to evaluate the following definite integral.

$$\int_0^{\pi/12} \cos 2x \, dx$$

³ Click to view the table of general integration formulas.

$$\int_0^{\pi/12} \cos 2x \, dx = \boxed{} \text{ (Type an exact answer.)}$$

3: General Integration Formulas

$$\int \cos ax \, dx = \frac{1}{a} \sin ax + C$$

$$\int \sin ax \, dx = -\frac{1}{a} \cos ax + C$$

$$\int \sec^2 ax \, dx = \frac{1}{a} \tan ax + C$$

$$\int \csc^2 ax \, dx = -\frac{1}{a} \cot ax + C$$

$$\int \sec ax \tan ax \, dx = \frac{1}{a} \sec ax + C$$

$$\int \csc ax \cot ax \, dx = -\frac{1}{a} \csc ax + C$$

$$\int e^{ax} \, dx = \frac{1}{a} e^{ax} + C$$

$$\int b^x \, dx = \frac{1}{\ln b} b^x + C, b > 0, b \neq 1$$

$$\int \frac{dx}{a^2 + x^2} = \frac{1}{a} \tan^{-1} \frac{x}{a} + C$$

$$\int \frac{dx}{\sqrt{a^2 - x^2}} = \sin^{-1} \frac{x}{a} + C, a > 0$$

$$\int \frac{dx}{x\sqrt{x^2 - a^2}} = \frac{1}{a} \sec^{-1} \left| \frac{x}{a} \right| + C, a > 0$$

Answer: $\frac{1}{4}$

ID: 5.5.45

138. Use a change of variables or the table to evaluate the following definite integral.

$$\int_0^1 6e^{3x} dx$$

⁴ Click to view the table of general integration formulas.

$$\int_0^1 6e^{3x} dx = \boxed{} \text{ (Type an exact answer.)}$$

4: General Integration Formulas

$$\int \cos ax \, dx = \frac{1}{a} \sin ax + C$$

$$\int \sin ax \, dx = -\frac{1}{a} \cos ax + C$$

$$\int \sec^2 ax \, dx = \frac{1}{a} \tan ax + C$$

$$\int \csc^2 ax \, dx = -\frac{1}{a} \cot ax + C$$

$$\int \sec ax \tan ax \, dx = \frac{1}{a} \sec ax + C$$

$$\int \csc ax \cot ax \, dx = -\frac{1}{a} \csc ax + C$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax} + C$$

$$\int b^x dx = \frac{1}{\ln b} b^x + C, b > 0, b \neq 1$$

$$\int \frac{dx}{a^2 + x^2} = \frac{1}{a} \tan^{-1} \frac{x}{a} + C$$

$$\int \frac{dx}{\sqrt{a^2 - x^2}} = \sin^{-1} \frac{x}{a} + C, a > 0$$

$$\int \frac{dx}{x\sqrt{x^2 - a^2}} = \frac{1}{a} \sec^{-1} \left| \frac{x}{a} \right| + C, a > 0$$

Answer: $2e^3 - 2$

ID: 5.5.46

139. Use a change of variables or the table to evaluate the following definite integral.

$$\int_0^1 4x^3 (9 - x^4) dx$$

⁵ Click to view the table of general integration formulas.

$$\int_0^1 4x^3 (9 - x^4) dx = \boxed{} \text{ (Type an exact answer.)}$$

5: General Integration Formulas

$$\int \cos ax \, dx = \frac{1}{a} \sin ax + C$$

$$\int \sin ax \, dx = -\frac{1}{a} \cos ax + C$$

$$\int \sec^2 ax \, dx = \frac{1}{a} \tan ax + C$$

$$\int \csc^2 ax \, dx = -\frac{1}{a} \cot ax + C$$

$$\int \sec ax \tan ax \, dx = \frac{1}{a} \sec ax + C$$

$$\int \csc ax \cot ax \, dx = -\frac{1}{a} \csc ax + C$$

$$\int e^{ax} \, dx = \frac{1}{a} e^{ax} + C$$

$$\int b^x \, dx = \frac{1}{\ln b} b^x + C, b > 0, b \neq 1$$

$$\int \frac{dx}{a^2 + x^2} = \frac{1}{a} \tan^{-1} \frac{x}{a} + C$$

$$\int \frac{dx}{\sqrt{a^2 - x^2}} = \sin^{-1} \frac{x}{a} + C, a > 0$$

$$\int \frac{dx}{x\sqrt{x^2 - a^2}} = \frac{1}{a} \sec^{-1} \left| \frac{x}{a} \right| + C, a > 0$$

Answer: $\frac{17}{2}$

ID: 5.5.47

140. Use a finite approximation to estimate the area under the graph of the given function on the stated interval as instructed.

$f(x) = x^2$ between $x = 0$ and $x = 2$, using a right sum with two rectangles of equal width

- ☐ A. 4.5
☐ B. 1
☐ C. 5
☐ D. 2.5

Answer: C. 5

ID: 5.1-1

141. Suppose that $\int_1^2 f(x)dx = -2$. Find $\int_1^2 3f(u)du$ and $\int_1^2 -f(u)du$.
-

- ☐ A. $-6; -\frac{1}{2}$
- ☐ B. $3; 2$
- ☐ C. $-6; 2$
- ☐ D. $1; -2$

Answer: C. $-6; 2$

ID: 5.2-12

142. Find the derivative.

$$\frac{d}{dx} \int_1^{\sqrt{x}} 16t^9 dt$$

- ☐ A. $\frac{16}{5}x^{3.5} - \frac{16}{5}$
- ☐ B. $16x^{9/2}$
- ☐ C. $\frac{32}{3}x^{3.5}$
- ☐ D. $8x^{1.5}4$

Answer: D. $8x^{1.5}4$

ID: 5.3-18

143. Evaluate the integral using the given substitution.

$$\int x \cos(10x^2) dx, \quad u = 10x^2$$

- ☐ A. $\frac{1}{u} \sin(u) + C$
- ☐ B. $\frac{x^2}{2} \sin(10x^2) + C$
- ☐ C. $\sin(10x^2) + C$
- ☐ D. $\frac{1}{20} \sin(10x^2) + C$

Answer: D. $\frac{1}{20} \sin(10x^2) + C$

ID: 5.5-1

144. Evaluate the following integral.

$$\int \frac{dx}{x^2 - 2x + 5}$$

$$\int \frac{dx}{x^2 - 2x + 5} = \boxed{}$$

(Use C as the arbitrary constant as needed.)

Answer: $\frac{1}{2} \tan^{-1} \frac{x-1}{2} + C$

ID: 8.1.31

145. If the general solution of a differential equation is $y(t) = C e^{-4t} + 10$, what is the solution that satisfies the initial condition $y(0) = 6$?

$$y(t) = \boxed{}$$

Answer: $-4 e^{-4t} + 10$

ID: 9.1.2

146. Find the general solution of the following equation. Express the solution explicitly as a function of the independent variable.

$$t^{-4}y'(t) = 4$$

$$y = \boxed{}$$

Answer: $\frac{4t^5}{5} + C$

ID: 9.3.5

147. Find the general solution of the differential equation $\frac{dy}{dt} = \frac{3t^2}{4y}$.

Choose the correct answer below.

- ☐ A. $y = 2t^3 + C$
- ☐ B. $y = \pm \sqrt{2t^3 + C}$
- ☐ C. $y = \pm \sqrt{\frac{t^3}{2} + C}$
- ☐ D. $y = \frac{t^3}{2} + C$

Answer: C. $y = \pm \sqrt{\frac{t^3}{2} + C}$

ID: 9.3.7

148. The general solution of a first-order linear differential equation is $y(t) = C e^{-10t} - 19$. What solution satisfies the initial condition $y(0) = 5$?

The solution is $y(t) = \boxed{}$.

Answer: $24 e^{-10t} - 19$

ID: 9.4.1

149. Find the general solution of the following equation.

$$y'(t) = 4y - 5$$

$$y(t) = \boxed{}$$

(Use C as the arbitrary constant.)

Answer: $C e^{4t} + \frac{5}{4}$

ID: 9.4.5

150. Find the general solution of the following equation.

$$y'(x) = -y + 4$$

$$y(x) = \boxed{}$$

(Use C as the arbitrary constant.)

Answer: $C e^{-x} + 4$

ID: 9.4.6